

# Rhodora

JOURNAL OF

THE NEW ENGLAND BOTANICAL CLUB

---

Vol. 9.

August, 1907.

No. 104.

---

## A PARTIAL LIST OF CONNECTICUT DIATOMS WITH SOME ACCOUNT OF THEIR DISTRIBUTION IN CERTAIN PARTS OF THE STATE.

WILLIAM A. TERRY.

CONNECTICUT is rich in diatoms, both in quantity and variety, and fossil deposits are abundant, both of fresh water and marine. The very ancient deposits like those of Virginia and Maryland cannot be expected here, as the great ice sheet during glacial times swept all such as may have existed into the sea, but the time since then has been sufficient for the accumulation of a vast store of material of which very little has ever been investigated. I have found eleven fresh water deposits in Bristol, varying in size from a few rods in diameter to many acres; one in the northern part of the town, "No. 11," covers fifteen acres and possibly much more. The diatomaceous stratum of these deposits is generally three to four feet below the surface, and in the larger deposits averages about two feet in depth, but in No. 11 it is of unknown depth; I have material from ten and a half feet below the surface, showing about seven feet of diatoms to this point; it is especially interesting as it contains numbers of the beautiful *Cyclotella antiqua* W. Sm., which is very rare in this country; all correspondents to whom I have sent it report that they had never seen it before. Dr. Ward sent me a slide with this label from Prof. H. L. Smith's collection which is in his possession, but the forms shown were not the same. The specimens from which Van Heurck made his drawings for his Synopsis must have been very inferior to the Bristol specimens in beauty.

In New Britain Mr. W. R. Stone discovered a fossil deposit in which I found a new *Stauroneis*, the outline of which resembled that of

*Navicula amphirhyncus* Ehrenb., but exhibited an oblique stauros. This might appropriately have been called *obliqua*, but as Gregory had appropriated this name for an entirely different and much smaller form, I named it *S. nova britannica*, and it was so published in Dr. Bigelow's *Observer*. Gregory's form was not strictly a *Stauroneis*; it was a very small stauroneiform *Scoliopleura*, and its stauros was not oblique. This New Britain *Stauroneis* is the most rare species of which I have any knowledge; Mr. Stone's original gathering contained some hundreds, perhaps thousands of specimens, but on subsequent investigation no more could be found; they were probably confined to a very thin stratum of the deposit which was not identified, and further exploration is difficult if not impossible as the ground is covered with buildings.

Connecticut waters also are rich in living diatoms, and recent deposits abound. Nearly every lake and pond has its bottom covered with a soft ooze, which is filled with diatoms and is often two or more feet in thickness. The margins of rivers, ditches in the marshes, and the bottoms of small streams and rills in springy mountain pastures, furnish them in abundance. A small stream in a marshy pasture on Fall Mountain in Bristol has an abundant colony of the large variety of *Stauroneis acuta* W. Sm., many times larger than the type of Wm. Smith. I have a slide sent me by Prof. H. L. Smith which he assured me was part of the original gathering from which Tuffen West made the drawing of *S. acuta* for Smith's *British Diatomaceae*; this is the same as the *Pleurotaenium acutum* of other authors, and is distinctly different from the large Bristol *Stauroneis*. In the colony I find another *Stauroneis* which is quite remarkable; it is longer and more cylindrical than the other species, has a broad stauros and rounded ends with a large saucer-shaped pseudonodule near each end of the upper valve but none on the lower valve. Next to the New Britain *Stauroneis* this is the most rare of all our species of diatoms. They may be gathered in hundreds from a space of about thirty feet on this brook, but they are so much outnumbered by other species with them that it is difficult to find them after they are collected; I have found a single specimen in each of three widely separated ponds in Bristol, but they are known nowhere else. Dr. Ward named this species *Stauroneis Terryi*. In another small stream on Fall Mountain, Mr. W. C. Richards found a colony of *Navicula elliptica* Kütz., notable for abundance and size of individuals.

In five different ponds in Bristol I have found an abundance of a new *Surirella*, and in one of these ponds it is the predominating form. It is about the size of *S. gracilis* A. Schm., but not so alate, and with rounded ends, costa distinct, reaching the median line, which is strongly marked; most specimens have a slight spiral twist. It occurs in two types, one greatly elongated. The living frustules are sometimes covered with coarse granulations that do not come off in the acid treatment, but are removed by the soda. Prof. H. L. Smith wrote me "This *Surirella* is certainly new and much more deserving of a specific name than many others." Dr. Ward named it *Surirella Terryi*. I have found this species abundant in one small pond in New Britain and in one at Leete's Island, but in all other localities known to me it is extremely rare. In Birge's Pond in Bristol are countless numbers of a very peculiar abnormal form of *Surirella*. It is a double frustule in which the two inner valves are grown together with a perforation through the center where the protoplasm of the two cells joins. This opening is sometimes a mere raphe or cleft, more often it is a long narrow slit with smooth edges, but generally it is a large irregular orifice with corrugated sides. This peculiarity is most abundant in *Surirella elegans* Ehrenb., but frequent in *S. cardinalis* Kitton, and occasional, in *S. splendida* (Ehrenb.) Kütz. I found it first in *S. striatula* Turpin in an artificial culture of material from West River, New Haven. Mr. Richards brought me mud from a pond in Wallingford, in which I found it abundant in *S. saxonica* Auersw.; I have seen it in *S. tenera* Greg. and *S. fastuosa* Ehrenb., and have found it in Plymouth and in Derby. An illustrated description was published in Tempère's *Micrographe Préparateur*.<sup>1</sup>

Nearly every marsh with a level surface covers a deposit of diatoms, and the salt marshes of the Connecticut shore are no exceptions. All of these that I have examined lie above a marine deposit, which generally appears like clay, but always contains diatoms and is sometimes quite rich. The ancient channel of Leete's Island from Great Harbor on the south around to Shell Beach on the north, is one immense mass of marine deposit, in some places fifty feet thick. When the railroad crossing it was changed from a trestle to an embankment, the pressure of the great bank of earth forced out the marine mud

<sup>1</sup> W. A. Terry, Sur un étrange mode de développement chez le genre *Surirella*. *Le Micrographe Préparateur*, Vol. XIII, p. 57, 1905.

below it which rose up in great banks on each side of the railroad. All this material contained diatoms, and some of it was very rich, especially in number and rarity of species. I dug up material from about twelve feet below the surface, and found a stratum rich in the beautiful *Surirella Febigerii* Lewis and other rare kinds. One pound of this earth furnished diatoms enough to make several thousand slides, each slide containing thousands of diatoms. The marine deposit in these marshes is sometimes near the surface, and is thrown out in digging ditches, and always in quantities in preparing the foundations for the abutments of bridges. From the earth thrown out in digging a post-hole for a fence on the margin of the marsh near Branford Station I procured a supply of *Campylodiscus echineis* Ehrenb. The Quinnipiac marshes are large, and along their western margin is a series of clay pits from which the tide water is kept out by dikes. The diatomaceous deposit here is of no great thickness but has a very interesting collection of species. At Davis pit the clay is covered by a layer of sand, upon which rests a stratum of alluvium, which contains fresh water diatoms exclusively; these are mostly decayed and broken, those entire being chiefly *Navicula lata* Bréb., with a few of the large *Stauroneis acuta* and *Navicula Semen* Ehrenb. Upon the alluvium is a mass of loam containing the roots and stumps of trees with their trunks and branches beside them, the whole covered with the marine deposit, which extends upward through about four feet of coarse peat.

At many places along the Connecticut shore the ancient marine deposit projects through the bank of sand forming the beach, and is visible at low tide, the huge bank of sand being a superficial structure cast up by the sea and resting upon the ancient deposit. All these salt marshes were once open water, and the sea is now coming back upon them and swallowing up the land. Some of the pond holes in these marshes are deep, with a bottom of soft mud; these are often the remains of former open water, and are always rich in diatoms, frequently very different from those of the neighboring waters. Here is the home of the large *Pleurosigma balticum*, the very large *Amphiprora pulchra* Bail., the larger type of *Surirella striatula*, *Scolio-pleura*, *Amphora*, and a variety of rare forms.

In a pond hole of the marshes of Morris Creek I found a new *Pleurosigma*, quite large, in Morris Creek another, not quite so large. On the rocks near the tide gate were masses of *Melosira Borreri* Grev.,

looking in the water like enormous sponges, but collapsing when gathered into a rope of harsh feeling filament, about two feet long, which held great numbers of other diatoms, among them another new *Pleurosigma*, quite small. I sent all three to Tempère, he passed them on to Peragallo, who published illustrated descriptions of them in his monograph<sup>1</sup> which is now the accepted standard for *Pleurosigma*. Peragallo named the largest one *P. Terryanum*, the second *P. Americanum*, and the third *P. paradoxum*. He describes the first as having the outline of *P. strigilis* W. Sm., but the striation of *P. balticum*; the second having the outline of *P. elongatum* W. Sm., but striation of *P. decorum* W. Sm.; and the third was named *P. paradoxum* because the least approximate of its series of striae was the most difficult to resolve. I afterwards found *P. Terryanum* more abundant and larger in a broad ditch at Quinnipiac, and in a pond at Leete's Island. Its natural habitat is in brackish water fed by fresh water springs and too nearly fresh to support *P. balticum*. With this is often found *P. elongatum*, and also the robust type of *Surirella striatula* and a peculiar variety of *Navicula permagna* (Bail.) Ehrenb. The normal type of *N. permagna* is abundant in Oyster River, Woodmont. The species of *Pleurosigma* are found with it at Leete's Island, and also a remarkable colony of *Navicula maculata* Bail., which occurs here in great abundance, and in three distinct types; the largest is extremely rare elsewhere, the second found occasionally along the Connecticut shore, and the third and smallest is the most common. I have it from the New Jersey shore and from Cuxhaven, North Sea. A much smaller variety but with coarser puncta is found in Alabama. The typical *P. Terryanum* is the largest *Pleurosigma* that I find in Connecticut; *P. Americanum* is abundant in tidal creeks; in the marsh on the West Haven side of West River, on the muddy sloping bank of a ditch, I have seen at low tide a brown film about two feet wide and a hundred feet long, which consisted almost entirely of living frustules of *P. Americanum*. *P. paradoxum* is most abundant in Morris Creek, occasional elsewhere. Morris Creek is also the habitat of *Actinocyclus Barkleyi* (Ehrenb.) Grun., which is living here in abundance. I have it also in considerable numbers from the Thames river above Norwich in material sent me by Mr. G. R. Lumsden of Greenville.

<sup>1</sup> Peragallo, Monographie du genre *Pleurosigma*. Le Diatomiste, Vol. I.

At Branford I find *Ditylum Brightwellii* (West) Grun.; the drawings in books show this with long central spine with abrupt termination, without finish; the Branford specimens have a longer spine terminating in a knob or ball, which indicates that the specimens from which the figures were made were broken. *Nitzschia scalaris* (Ehrenb.) W. Sm. grows abundantly in the brackish water of many of the pond holes of the salt marshes. Pond's Point at Milford is a rounded projection where the southward trend of the shore changes abruptly to west. The beach is composed of large cobblestones and boulders upon a coarse gravel. On its southern margin is visible at low tide a bank of fossil mud, which is firm enough to resist the action of the waves, and is being gradually uncovered as the mass of stones is driven back by storms. A stratum near the upper part of this bank is very rich in diatoms, showing an unusual combination of marine and brackish water forms of more than one hundred species. *Coscinodiscus*, *Actinoptychus*, *Triceratium*, *Amphitetras*, *Eupodiscus*, with a dozen species of *Pleurosigma* occur, while the abundance of the brackish water form of *Surirella striatula* and *Navicula permagna*, with *Scolio-pleura*, *Amphora*, *Amphiprora* and *Nitzschia scalaris* (Ehrenb.) W. Sm. show this mud to have once formed the bottom of a pond hole in the marsh; the fresh water spring that fed it still flows over it, and can be seen at low tide. The pressure of the great bank of stones which rested upon it during the time it was being driven over by the sea has crushed to fragments most of the delicate species, but enough remain to show conclusively its character. The marine species mentioned above can be found generally in deep water all along the Connecticut shore, while the different varieties of *Navicula lyra* Ehrenb. and the *Pleurosigmas* are most abundant in shallow water. I have *P. intermedium* W. Sm. abundant in the deep water of Clinton, and the very rare *Navicula Lewisiana* Grev. in shallow water at Bridgeport. *P. balticum* is the most abundant species of the genus, and has many varieties, the largest being found in the salt marshes, the normal type in coves and bays, and the smallest in deep water. *P. affine* Grun. is the most universally distributed being found everywhere along the shore but not generally in great abundance. It is very often mistaken for *P. angulatum* W. Sm., although striation is very different. *P. angulatum* is frequent but local. A ditch in the Quinnipiac marshes through which flows at low tide a strong stream of nearly fresh water is carpeted with a dense growth of *Biddulphia levis* Ehrenb., and near

this is a colony of large and very active *Bacillaria paradoxa* (Gmel.) Grun. Another colony of this species in Morris Creek marsh, although in constant motion, was covered with a parasitic growth of *Leptothrix*, and was accompanied by numerous rapidly revolving filaments of *Spirulina*.

In fresh water the most abundant genus is *Pinnularia* W. Sm.; it is found in quantities almost everywhere. Next to this in numbers is the genus *Stauroneis*, then *Cymbella*, *Surirella*, *Navicula*, and numerous filamentous forms. Although these genera are represented in nearly every pond, yet each pond shows a combination characteristic of itself, and types of the same species vary greatly in the different ponds. In several ponds in Bristol *Surirella elegans* is very large and sometimes much elongated; in Plymouth it is smaller and much shorter in proportion; in Highland Lake, Winsted, beautifully typical specimens of *S. nobilis* W. Sm. are abundant, and in several ponds in Connecticut *Navicula peripunctata* Brun abounds; Prof. Brun founded the species on specimens in mud from Crane Pond, Groveland, Massachusetts, but it is more plentiful in Bristol and in Leete's Island. Van Heurck states that *Achnanthisidium flexillum* (Kütz.) Bréb. is rare; it is abundant in two of the Bristol fossil deposits. He also gives *Fragilaria Harrisonii* Grun as very rare; this grows abundantly in Bristol. Dr. Ward found a new diatom in one of my slides from the Connecticut shore, which Cleve named *Caloneis Wardii*. Mr. O. E. Shaffer sent a find of *Nitzschia* to Europe from Port Townsend, Puget Sound, and Cleve named this *N. Shafferi*; I had previously found this nearly twenty years ago at Morris Creek, and reported it at the time in the Amer. Micr. Journal as *N. curvula*. I find *Isthmia nervosa* Kütz. in nearly every deep water sounding from off the Connecticut shore, but do not believe that it grows here; it is probably brought in by ocean currents from warmer waters.

#### LIST OF SPECIES.

The arrangement of this list is that of De Toni, *Sylloge Algarum*, and the nomenclature for the most part follows that work. Names given as synonyms are those that have been used by the writer either in published notes or with material distributed. Species marked with a star were founded on Connecticut material.

*Navicula nobilis* (Ehrenb.) Kütz. Common.

- Navicula* var. *dactylus* (Ehrenb.) VH. (*N. dactylus* Kütz.) Abundant everywhere.
- “ *gigas* (Ehrenb.) Kütz. Common.
- “ *major* Kütz. Abundant everywhere.
- “ *viridis* (Nitzsch) Kütz. Abundant everywhere.
- “ *cardinalis* Ehrenb. Frequent, local, Bristol.
- “ *yarrensis* Grun. Occasional.
- “ *divergens* (W. Sm.) Ralfs. Frequent, local, Bristol.
- “ *longa* (Greg.) Ralfs. Frequent.
- “ *lata* Bréb. Frequent.
- “ *borealis* (Ehrenb.) Kütz. Frequent, local, Bristol.
- “ *directa* Ralfs. Occasional.
- “ *Brebissonii* Kütz. Frequent, local, Bristol.
- “ *Tabellaria* Kütz. Frequent, local.
- “ *gibba* (Ehrenb.) Kütz. Frequent, local, Bristol.
- “ *Legumen* Ehrenb. Occasional.
- “ *polyonca* Bréb. Frequent.
- “ *peregrina* (Ehrenb.) Kütz. Common.
- “ *semen* Ehrenb. Frequent, local, Bristol.
- “ *zostereti* Grun. Rare, Morris Creek.
- “ *rhyncocephala* Kütz. Occasional.
- “ *cancellata* Donk. Frequent, local, Fort Hale.
- “ *fortis* (Greg.) Grun. Occasional.
- “ *digito-radiata* (Greg.) Ralfs. Occasional.
- “ *distans* (W. Sm.) Ralfs. Occasional.
- “ *brasiliensis* Grun. Frequent.
- “ *crabro* (Ehrenb.) Kütz. Frequent.
- “ *interrupta* (Bail.) Kütz. Frequent.
- “ *didyma* Ehrenb. Common.
- “ *bombus* (Ehrenb.) Kütz. Occasional.
- “ *Smithii* Bréb. Frequent.
- “ *fusca* (Greg.) Ralfs. Frequent.
- “ *advena* A. Schm. Occasional.
- “ *elliptica* Kütz. Occasional.
- “ “ var. *oblongella* (Näg.) VH. (*N. oblongella* Näg.) Occasional.
- “ *lyra* Ehrenb. Locally abundant.
- “ *pygmaea* Kütz. Local.
- “ *forcipata* Grev. Occasional, New Haven Harbor.
- “ *caribaea* Cleve. Occasional, New Haven Harbor.
- “ *approximata* Grev. Occasional, New Haven Harbor.
- “ *irrorata* Grev. Occasional.
- “ *praetexta* Ehrenb. Occasional.
- “ *Hennedyi* W. Sm. Occasional, Leete's Island.
- “ *clavata* Greg. Occasional.
- “ *polysticta* Grev. Occasional, New Haven Harbor.
- “ *hibernica* O'M. Occasional.

- Navicula aspera* Ehrenb. Occasional.  
 “ *californica* var. *campeachiana* Grun. Occasional.  
 “ *circumsecta* Grun. Occasional.  
 “ *tuscula* Ehrenb. Rare, local, Bristol.  
 “ *crucicola* (W. Sm.) Donk. Local.  
 “ *palpebralis* Bréb. Occasional.  
 “ *Fischeri* A. Schm. Occasional.  
 “ *arabica* Grun. Occasional.  
 “ *humerosa* Bréb. Frequent.  
 “ *tumescens* Grun. Common.  
 “ *marina* Ralfs. Occasional.  
 “ *pusilla* W. Sm. Local, Quinnipiac.  
 “ *latissima* Greg. Rare.  
 “ *kamorthensis* Grun. Rare, Stony Creek.  
 “ *cuspidata* Kütz. Frequent, local, Bristol.  
 “ *sphaerophora* Kütz. Rare, local, Bristol.  
 “ *serians* (Bréb.) Kütz. Frequent, local, Bristol.  
 “ *formosa* Greg. Frequent.  
 “ *permagna* (Bail.) Edw. Occasional; abundant at Woodmont.  
 “ *elegans* W. Sm. Occasional.  
 “ *limosa* Kütz. Occasional.  
 “ *iridis* Ehrenb. Common.  
 “ “ var. *Amphigomphus* (Ehrenb.) VH. (*N. amphigomphus* Ehrenb.) Local.  
 “ “ var. *producta* (W. Sm.) VH. (*N. producta* W. Sm.) Occasional.  
 “ “ var. *amphirhynchus* (Ehrenb.) De Toni. (*N. amphirhynchus* Ehrenb.) Local.  
 “ *firma* Kütz. Common.  
 “ *Hitchcockii* Ehrenb. Rare, local, Bristol.  
 “ *liber* W. Sm. Occasional.  
 “ *americana* Ehrenb. Rare, local, Bristol.  
 “ *bacillum* Ehrenb. Local.  
 “ *Powellii* Lewis. Occasional.  
 “ *columnaris* Ehrenb. Common.  
 “ *esox* (Ehrenb.) Kütz. Occasional.  
 “ *Sillimanorum* (Ehrenb.) Kütz. Rare, local, Bristol.  
 “ *\*parallela* Castr. Rare, Bristol.  
 “ *dilatata* Ehrenb. Common.  
 “ *delawarensis* Grun. Local, Quinnipiac.  
 “ *interposita* Lewis. Frequent.  
 “ *amphipleuroides* Grun. Rare.  
 “ *Bartleyana* Grun. Occasional.  
 “ *diploneis* var. *obliquus* Brun. Occasional.  
 “ *maculata* Bail. Occasional; abundant at Leete’s Island.  
 “ *peripunctata* Brun. Occasional; abundant in Bristol.

- Navicula* \**theta* Cleve. Local, Quinnipiac.  
 " \**tubulosa* Brun. Occasional.  
*Caloneis* \**Wardii* Cleve. Occasional.  
*Diploneis* \**didyma* var. *obliqua* Brun. Occasional.  
*Libellus rhombicus* (Greg.) De Toni. (*Navicula rhombica* Greg.)  
*Rhoiconeis trinodis* (W. Sm.) Grev. (*Navicula trinodis* W. Sm.)  
 Rare, Bristol.  
*Stauroneis phoenicenteron* (Nitzsch) Ehrenb. Abundant everywhere.  
 " *gracilis* W. Sm. Frequent.  
 " *Gregoryi* Ralfs. Occasional.  
 " *spicula* Hickie. Rare, Quinnipiac.  
 " *salina* W. Sm. Common along shore.  
 " *Stodderi* Lewis. Rare, Bristol.  
 " \**nova britannica* Terry. Rare; only in New Britain.  
 " \**Terryi* Ward. Exceedingly rare; only in Bristol.  
*Pleurostauron acutum* (W. Sm.) Ralfs. (*Stauroneis acuta* W. Sm.)  
 Common.  
*Pleurosigma angulatum* (Quek.) W. Sm. Frequent, local.  
 " " var. *aestuarii* (Bréb.) VH. Occasional.  
 " *elongatum* W. Sm. Frequent.  
 " " var. *gracile* Grun. Occasional.  
 " *strigosum* W. Sm. Rare, Clinton.  
 " *naviculaceum* Bréb. Rare.  
 " *intermedium* W. Sm. Rare, Clinton.  
 " *nubecula* W. Sm. Rare, Clinton,  
 " *latum* Cleve. Rare.  
 " *formosum* W. Sm. Occasional.  
 " *hypocampus* W. Sm. Frequent, local.  
 " *decorum* W. Sm. Locally abundant.  
 " *littorale* W. Sm. Rather rare.  
 " *balticum* (Ehrenb.) W. Sm. Frequent, local.  
 " *simile* Grun. Rare, Leete's Island.  
 " *attenuatum* W. Sm. Frequent, local.  
 " *Wansbeckii* Donk. Frequent.  
 " *acuminatum* (Kütz.) Grun. Rather rare.  
 " *strigile* W. Sm. Occasional.  
 " *affine* Grun. & Cleve. Common all along the shore.  
 " \**paradoxum* Per. Occasional, Morris Creek.  
 " *Brebissonii* Grun. & Cleve. Frequent.  
 " \**Terryanum* Per. Local, Quinnipiac, Leete's Island.  
 " *Spencerii* (Quek.) W. Sm. Frequent, local.  
 " " var. *curvulum* (Kütz.) Grun. Rare, Quinnipiac.  
 " *fasciola* W. Sm. Frequent.  
 " *scalpoides* Rab. Rare.  
 " *eximium* (Thw.) Grun. & Cleve. Occasional.  
 " \**americanum* Per. Abundant, Morris Creek, New Haven.

- Pleurosigma subsalinum* Per. Rare, Leete's Island, Milford.  
*Gyrosigma* \**Temperii* Cleve. Occasional.  
*Scoliopleura latestriata* (Bréb.) Grun. Frequent.  
 " " \*var. *Amphora* Cleve. Rare, Morris Creek.  
 " *tumida* (Bréb.) Rab. Frequent.  
*Frustulia rhomboides* (Ehrenb.) De Toni. (*Navicula rhomboides* Ehrenb. Frequent.  
 " " var. *saxonica* (Rab.) De Toni. (*Navicula crassinervia* Bréb.) Occasional.  
 " *Lewisiana* (Grev.) De Toni. (*Navicula Lewisiana* Grev.) Rare, Bridgeport only.  
 " *viridula* (Bréb.) De Toni. (*Schizonema viridulum* Rab.) Rare.  
*Dickiea crucigera* (W. Sm.) De Toni. (*Schizonema crucigerum* W. Sm.) Rare, Quinnipiac.  
*Brebissonia Boeckii* (Kütz.) Grun. (*Navicula Boeckii* Bréb.) Occasional; abundant in Thames River.  
*Mastogloia* \**gibbosa* Brun. Occasional.  
 " *undulata* Grun. Occasional.  
*Amphiprora lepidoptera* Greg. Frequent.  
 " " var. *pusilla* (Greg.) VH. Frequent.  
 " *maxima* Greg. Locally abundant.  
 " *alata* Kütz. Frequent.  
 " *pulchra* Bail. Locally abundant.  
 " *conspicua* Grev. Frequent.  
 " *ornata* Bail. Frequent.  
*Plagiotropis vitrea* (W. Sm.) Grun. (*Amphiprora vitrea* W. Sm.) Frequent.  
*Tropidoneis* \**seriata* Cleve. Occasional.  
 " \**zebra* Cleve. Occasional.  
*Cymbella Ehrenbergii* Kütz. Locally abundant.  
 " *cuspidata* Kütz. Frequent.  
 " *affinis* Kütz. Local, New Britain.  
 " *heteropleura* (Ehrenb.) Kütz. Local.  
 " *gastroides* Kütz. Common everywhere.  
 " *cymbiformis* var. *parva* (W. Sm.) VH. Local, New Britain.  
 " *americana* A. Schm. Locally abundant.  
 " *cistula* var. *maculata* (Kütz.) Grun. Local, New Britain.  
*Encyonema caespitosum* Kütz. Frequent.  
 " *prostratum* (Bail.) Ralfs. Frequent.  
*Amphora levis* Greg. Frequent.  
 " *cingulata* Cleve. Frequent.  
 " *erebi* Ehrenb. (*A. cymbifera* Greg.) Frequent.  
 " *Clevei* A. Schm. Frequent.  
 " *intersecta* A. Schm. Frequent.  
 " *Eulensteinii* Grun. Frequent; locally abundant at Milford.  
 " *proteus* Greg. Frequent.

- Amphora obtusa* Greg. Frequent.  
 “ *ovalis* (Bréb.) Kütz. Frequent.  
*Gomphonema acuminatum* Ehrenb. Common.  
 “ *constrictum* Ehrenb. Common.  
 “ *capitatum* Ehrenb. Common.  
 “ *geminatum* (Lyng.) Ehrenb. Common.  
 “ *augur* Ehrenb. (*G. cristatum* Ralfs.) Common.  
*Rhoicosphaenia curvata* (Kütz.) Grun.  
*Cocconeis scutellum* Ehrenb. Frequent.  
 “ *dirupta* Greg. Frequent.  
*Achnanthes longipes* Ag. Common.  
 “ *brevipes* Ag. Common.  
 “ *subsessilis* Kütz. Fort Hale.  
 “ *\*curvirostrum* Brun. Very rare, Morris Cove.  
 “ *\*manifera* Brun. Very rare, Morris Cove.  
*Achnanthidium flexellum* (Kütz.) Bréb. Rare; abundant in Bristol,  
 fossil.  
*Bacillaria paradoxa* (Gmel.) Grev. Occasional near New Haven.  
*Nitzschia granulata* Grun. Occasional.  
 “ *tryblionella* Hantzsch. Frequent.  
 “ “ var. *maxima* Grun. Occasional.  
 “ *plana* W. Sm., Occasional.  
 “ *marginulata* Grun. Occasional.  
 “ *acuminata* (W. Sm.) Grun. Occasional.  
 “ *circumsuta* (Bail.) Grun. Frequent.  
 “ *scalaris* (Ehrenb.) W. Sm. Abundant.  
 “ *angularis* W. Sm. Frequent.  
 “ *sigmoidea* (Nitzsch) W. Sm. Frequent, locally abundant.  
 “ *sigma* (Kütz.) W. Sm. Common.  
 “ *fasciculata* Grun. Common.  
 “ *obtusa* W. Sm. Occasional.  
 “ *linearis* var. *tenuis* (W. Sm.) Grun. (*N. tenuis* W. Sm.)  
 Rare, Leete's Island.  
 “ *curvirostris* var. *Closterium* (Ehrenb.) VH. (*N. Closterium*  
 W. Sm.) Rare, West Haven.  
 “ *Lorenziana* Grun. Rare.  
 “ *Shafferi* Cleve. (*N. curvula* Terry, not W. Sm.) Very  
 rare, Morris Cove.  
*Hantzschia amphioxys* (Ehrenb.) Grun. Abundant in Bristol.  
 “ *\*segmentalis* Brun. Rare, Morris Cove.  
*Surirella biseriata* (Ehrenb.) Bréb. Common.  
 “ *linearis* W. Sm. Frequent.  
 “ *robusta* Ehrenb. Frequent in Winsted.  
 “ *splendida* (Ehrenb.) Kütz. Common.  
 “ *tenera* Greg. Common.  
 “ *elegans* Ehrenb. Common.  
 “ “ var. *norvegica* (Eulen.) Brun. Occasional, Bristol.

- Surirella striatula* Turp. Frequent, locally abundant.  
 “ “ var. *triplicata* Grun. Occasional.  
 “ *gemma* Ehrenb. Frequent.  
 “ *cardinalis* Kitt. Rather rare; abundant in Bristol.  
 “ *ovalis* Bréb. Occasional.  
 “ “ var. *angusta* (Kütz.) VH. (*S. angusta* Kütz.) Occasional.  
 “ *fastuosa* Ehrenb. Frequent.  
 “ “ var. *lata* (W. Sm.) VH. Frequent.  
 “ *recedens* A. Schm. Occasional.  
 “ *Moelleriana* Grun. Occasional.  
 “ *gracilis* A. Schm. Occasional.  
 “ *Febigerii* Lewis. Frequent.  
 “ *constricta* Ehrenb. Rare, Killingsworth.  
 “ \**Terryi* Ward. Rare, except in Bristol, New Britain and Leete’s Island.
- Cymatopleura elliptica* (Bréb.) W. Sm. Rare, New Britain.  
 “ *solea* (Bréb.) W. Sm. Occasional.  
 “ *hibernica* W. Sm. Rare.  
 “ *marina* Lewis. Rare, Clinton.
- Podocystis adriatica* Kütz. (*P. americana* Bail.) Rare.
- Campylodiscus echineis* Ehrenb. Common.  
 “ *hibernicus* Ehrenb. Frequent.
- Diatoma vulgare* Bory. Frequent.  
 “ *elongatum* var. *tenue* (Ag.) VH. (*D. tenue* Ag.) Frequent.  
 “ *hiemale* (Lyngb.) Heib. (*Odontidium hiemale* Kütz.) Locally abundant.  
 “ *anceps* (Ehrenb.) Kirchn. Frequent.
- Odontidium mutabile* W. Sm. Locally abundant.  
 “ *Harrisonii* W. Sm. (*Fragilaria Harrisonii* Grun.) Rare and local, Bristol.
- Meridion circulare* (Grev.) Ag. Frequent.  
 “ *intermedium* H. L. Sm. Frequent.
- Synedra pulchella* (Ralfs) Kütz. Occasional.  
 “ “ var. *Smithii* (Ralfs) VH. Occasional.  
 “ *ulna* var. *danica* (Kütz.) VH. (*S. danica* Kütz.) Frequent.  
 “ *affinis* Kütz. Frequent.
- Ardissonia superba* (Kütz.) Grun. (*Synedra superba* Kütz.) Occasional, Leete’s Island.
- Asterionella formosa* Hass. Abundant in water supply.  
 “ *Ralfsii* W. Sm. Abundant in water supply.
- Fragilaria virescens* Ralfs. Common.  
 “ *capucina* Desmaz. Common.  
 “ *construens* (Ehrenb.) Grun. Common.
- Raphoneis amphiceros* Ehrenb. Frequent.  
 “ *belgica* Grun. Rare, Leete’s Island.  
 “ “ var. *intermedia* Grun. Occasional.  
 “ *gemmaifera* Ehrenb. Frequent.

- Dimerogramma minus* (Greg.) Ralfs.  
 “ *Surirella* (Ehrenb.) Grun. (*Raphoneis Surirella* Grev.) Occasional.
- Plagiogramma staurophorum* (Greg.) Heib. (*P. Gregorianum* Grev.) Occasional, Quinipiac.  
 “ *validum* Grev. Occasional, Leete’s Island.
- Licmophora flabellata* (Carm.) Ag. Frequent.  
 “ *Juergensii* Ag. Frequent.  
 “ *tincta* (Ag.) Grun. Frequent.
- Tabellaria fenestrata* (Lyngb.) Kütz. Common.  
 “ *flocculosa* (Roth) Kütz. Common.
- Grammatophora oceanica* var. *macilenta* (W. Sm.) Grun.  
*Rhabdonema arcuatum* (Lyngb.) Kütz. Occasional.  
 “ *adriaticum* Kütz. Common.
- Epithemia turgida* (Ehrenb.) Kütz. Frequent.  
 “ *gibba* (Ehrenb.) Kütz. Occasional.  
 “ “ var. *ventricosa* (Ehrenb.) Grun. Frequent.  
 “ *argus* (Ehrenb.) Kütz. Frequent.  
 “ *zebra* (Ehrenb.) Kütz. Frequent.  
 “ *musculus* Kütz. Frequent.
- Eunotia major* (W. Sm.) Rab. Frequent.  
 “ *praerupta* Ehrenb. Frequent.  
 “ “ var. *bidens* Grun. Frequent.  
 “ *gracilis* (Ehrenb.) Rab. Frequent.  
 “ *robusta* Ralfs, including *E. Diadema* Ralfs. Frequent.  
 “ *elongata* Rab. Frequent.
- Pseudoeunotia lunaris* (Ehrenb.) Grun. (*Eunotia lunaris* Grun.) Locally abundant.  
 “ *flexuosa* (Bréb.) Grun. (*Eunotia flexuosa* Bréb.) Locally abundant.  
 “ *flexuosa* var. *bicapitata* Grun. Locally abundant.
- Actinella punctata* Lewis. Common.  
*Rhizosolenia styliiformis* Brightw. Leete’s Island.  
*Isthmia nervosa* Kütz. Frequent.  
*Hemiaulus affinis* Grun. South Haven.  
 “ *tubulosus* Brun. Occasional.
- Odontella aurita* (Lyng.) Ag. (*Biddulphia aurita* Bréb.) Frequent.  
 “ *turgida* (W. Sm.) VH. Frequent.
- Biddulphia pulchella* Gray. Frequent.  
 “ *Tuomeyi* (Bail.) Roper. Occasional.
- Denticella mobiliensis* (Bail.) Grev. (*Biddulphia Baileyi* W. Sm.) Rare.  
 “ *rhombus* Ehrenb. (*Biddulphia rhombus* W. Sm.) Frequent.
- Eunotogramma laeve* Grun. Occasional.  
*Terpsinoe musica* Ehrenb. Occasional.  
 “ *americana* (Bail.) Ralfs. Occasional.

- Amphitetras antediluvianus* Ehrenb. Frequent.  
*Triceratium alternans* Bail. Occasional.  
 “ *flavum* Ehrenb. Frequent.  
*Lithodesmium undulatum* Ehrenb. Leete's Island.  
*Syndendrium diadema* Ehrenb. Occasional.  
*Periptera tetracladia* Ehrenb. Occasional.  
*Pyxilla baltica* Grun. Frequent; locally abundant, Leete's Island.  
*Ditylium Brightwellii* (West) Grun. including *D. inaequale* Bail.,  
*D. tetragonum* Bail., and *D. trigonum* Bail. Very rare,  
 Branford only.  
 “ *intricatum* (West) Grun. Very rare, Branford.  
*Auliscus pruinosis* Bail. Frequent.  
 “ *sculptus* (W. Sm.) Ralfs. Common.  
 “ *caelatus* Bail. Occasional.  
 “ *Macraëanus* Grev. Occasional.  
*Pseudoauliscus radiatus* (Bail.) Rattray. Rare, Leete's Island.  
*Cerataulus levis* (Ehrenb.) Ralfs. (*C. polymorphus* VH. *Biddulphia*  
*levis* Ehrenb.) Frequent.  
*Eupodiscus Rodgersii* Ehrenb. Rare, Leete's Island.  
 “ *argus* Ehrenb. Frequent.  
*Stephanopyxis turris* (Grev.) Ralfs. Occasional, New Haven.  
 “ “ *var. intermedia* Grun. (*S. appendiculata var. inter-*  
*media* Grun.) Occasional.  
 “ *ferox* (Grev.) Ralfs. Occasional, Milford.  
 “ *corona* (Ehrenb.) Grun. Occasional.  
*Stephanodiscus niagarae* Ehrenb. Rare.  
*Actinocyclus crassus* VH. Frequent.  
 “ *Ralfsii* (W. Sm.) Ralfs. Frequent.  
 “ *Barkleyi* (Ehrenb.) Grun. Abundant, Morris Cove.  
 “ *Ehrenbergii* Ralfs. Frequent.  
 “ *tenuissimus* Cleve. Rare, Morris Cove.  
 “ *subtilis* (Greg.) Ralfs. Rare, Morris Cove.  
 “ *triradiatus* Roper. Rare, Leete's Island.  
*Coscinodiscus excentricus* Ehrenb. Frequent.  
 “ *nitidus* Greg. Frequent.  
 “ *symbolophorus* Grun. Occasional.  
 “ *subtilis* Ehrenb. Rare, local.  
 “ *marginatus* Ehrenb. Occasional.  
 “ *radiatus* Ehrenb. Frequent.  
 “ *concinus* W. Sm. Frequent.  
 “ *gigas* Ehrenb. Rare, Leete's Island.  
 “ *asteromphalus* Ehrenb. Frequent.  
 “ *oculus-iridis* Ehrenb. Frequent.  
 “ *apiculatus* Ehrenb. Frequent.  
*Lysigonium moniliforme* (Mull.) Link. (*Melosira Borreri* Grev.)  
 Locally abundant.  
 “ *varians* (Ag.) De Toni. (*Melosira varians* Ag.) Abun-  
 dant.

- Gallionella nummuloides* (Dillw.) Bory. (*Melosira nummuloides* Ag.)  
Frequent.
- Melosira undulata* (Ehrenb.) Kütz. Frequent.
- “ *sculpta* (Ehrenb.) Kütz. Common.
- “ *coronata* Grun. New Haven.
- “ *octogona* Grun. Frequent.
- Paralia sulcata* (Ehrenb.) Cleve. Rare, Leete's Island.
- Cyclotella antiqua* W. Sm. Very rare, Bristol only.
- “ *striata* (Kütz.) Grun. Silver Sands.
- “ *Kuetzingiana* Thw. Frequent.
- Podosira dubia* (Kütz.) Grun. Occasional.
- Hyalodiscus subtilis* Bail. Occasional.
- “ *stelliger* Bail. Occasional.
- Actinoptychus undulatus* (Bail.) Ralfs. Common.
- “ *velatus* Ehrenb. Occasional.

BRISTOL, CONNECTICUT.

## THE GENUS *SUAEDA* IN NORTHEASTERN AMERICA.

M. L. FERNALD.

THE genus *Suaeda* has long been for the American botanist a source of much confusion and difficulty, in part because it consists of unattractive plants of saline soils which are generally ignored by collectors, in part because of the very indefinite conception of the true *Suaeda maritima* of Europe — the species about which our studies, at least of the eastern coast forms, must largely center. It has long been known to many New England botanists that we have on our northeastern coast more than the single species allotted to this region in Gray's Manual; and the two species described and figured (as *Dondia*) in Britton & Brown's Illustrated Flora satisfactorily cover only the forms which occur on the salt marshes and sea beaches from southern New England southward. On the coast of Maine are other forms which in habit, flowering-season, and fruit-characters are unlike the plants of the more southern shores.

During the summer of 1898 Mr. J. C. Parlin and the writer spent a day (July 23) upon Wells Beach and the adjacent marshes in southern Maine. At that time a small depressed *Suaeda* (no. 1) with short subcylindric dark green leaves was in mature fruit. Another depressed